arXiv:1410.5252v1 [math.CV] 20 Oct 2014

CRITERIA FOR UNIVALENCE AND QUASICONFORMAL EXTENSION OF HARMONIC MAPPINGS IN TERMS OF THE SCHWARZIAN DERIVATIVE

RODRIGO HERNÁNDEZ AND MARÍA J. MARTÍN

Abstract. We prove that if the Schwarzian norm of a given complex-valued locally univalent harmonic mapping f in the unit disk is small enough, then f is, indeed, globally univalent and can be extended to a quasiconformal mapping in the extended complex plane.

Introduction

In 1949, Nehari [9] proved that if a locally univalent analytic function φ in the unit disk $\mathbb{D}$ satisfies

$$\sup_{z\in\mathbb{D}} |S(\varphi)(z)|\,(1-|z|^2)^2 \leq 2\,,$$

then φ is globally univalent in $\mathbb{D}$. Here, $S(\varphi)$ denotes the *Schwarzian derivative* of φ defined by

$$S(\varphi) = \left(\frac{\varphi''}{\varphi'}\right)' - \frac{1}{2}\left(\frac{\varphi''}{\varphi'}\right)^2 . \tag{1}$$

Ahlfors and Weill [1] generalized Nehari's criterion of univalence by proving that if such function φ satisfies $\|S(\varphi)\| \leq 2t$ for some $t < 1$, then φ is injective in $\mathbb{D}$ and has a K-quasiconformal extension to $\widehat{\mathbb{C}} = \mathbb{C}\cup\{\infty\}$, where $K = (1+t)/(1-t)$.

Let now f be a complex-valued locally univalent harmonic mapping in the unit disk. By considering the complex conjugate, if needed, we can assume that f is sense-preserving. This is, $f = h + \overline{g}$ where h and g are analytic functions in $\mathbb{D}$ such that h is locally univalent and the

Date: October 21, 2014.

2010 *Mathematics Subject Classification.* 31A05, 30C55, 30C62.

Key words and phrases. Harmonic mappings, Schwarzian derivative, univalence criterion, quasiconformal extension.

This research was supported by grants Fondecyt 1110160 and 1110321, Chile. The second author is also supported by Academy of Finland grant 268009 and by Spanish MINECO Research Project MTM2012-37436-C02-02.

(second complex) dilatation $\omega = g'/h'$ is an analytic function mapping the unit disk into itself. The following definition for the Schwarzian derivative of such functions f was presented in [5]:

$$S_f = S(h) + \frac{\overline{\omega}}{1-|\omega|^2}\left(\frac{h''}{h'}\,\omega' - \omega''\right) - \frac{3}{2}\left(\frac{\overline{\omega}\,\omega'}{1-|\omega|^2}\right)^2, \tag{2}$$

where $S(h)$ is the classical Schwarzian derivative of h, as in (1).

The main purpose in this note is to prove that there exists a constant $\delta_0 > 0$ such that if the locally univalent harmonic mapping f in the unit disk has Schwarzian norm

$$\|S_f\| = \sup_{z\in\mathbb{D}} |S_f(z)|\,(1-|z|^2)^2 \le \delta_0\,,$$

then f is one-to-one in $\mathbb{D}$. We will also see that if $\|S_f\| \le \delta_0 t$ for some $t < 1$, then f has a quasiconformal extension to $\widehat{\mathbb{C}}$.

1. Background

1.1. **The Schwarzian derivative.** As it was mentioned in the introduction, every harmonic mapping f in the unit disk $\mathbb{D}$ can be written as $f = h + \overline{g}$ with h and g analytic in $\mathbb{D}$. This decomposition is unique up to an additive constant (see [3, p. 7]). We refer to the reader to the book [3] for a comprehensive treatment on harmonic mappings.

Lewy [8] proved that a harmonic mapping in the unit disk is locally univalent if and only if its Jacobian is different from zero. In terms of the decomposition $f = h + \overline{g}$, the Jacobian J_f of f equals $|h'|^2 - |g'|^2$. Thus, locally univalent harmonic mappings in $\mathbb{D}$ are either *sense-preserving* if $J_f > 0$ or *sense-reversing* if $J_f < 0$. Note that any analytic function is a sense-preserving harmonic mapping. Also, that a harmonic function $f = h+\overline{g}$ is sense preserving if and only if h is locally univalent and the dilatation $\omega = g'/h'$ maps the unit disk into itself. It is obvious that f is sense-preserving if and only if $\overline{f}$ is sense-reversing. In this paper, we will consider harmonic mappings which are sense-preserving in the unit disk. For this kind of mappings, the Schwarzian derivative is given by (2). It is clear that if f is analytic, then S_f coincides with the classical definition of the Schwarzian derivative given by (1).

Several properties of this operator are the following.

(i) $S_f \equiv 0$ if and only if $f = \alpha T + \beta\overline{T}$, where $|\alpha| \neq |\beta|$ and T is a Möbius transformation

$$T(z) = \frac{az+b}{cz+d}\,, \quad ad - bc \neq 0\,.$$

(ii) Whenever f is a sense-preserving harmonic mapping and ϕ is an analytic function such that the composition $f\circ\phi$ is well-defined, the Schwarzian derivative of $f\circ\phi$ can be computed using the *chain rule*

$$S_{f\circ\phi} = S_f(\phi)\cdot(\phi')^2 + S\phi\,.$$

(iii) For any *affine mapping* $L(z) = az + b\overline{z}$ with $|a| \neq |b|$, we have that $S_{L\circ f} = S_f$. Note that L is sense-preserving if and only if $|b| < |a|$.

The *Schwarzian norm* $\|S_f\|$ of a sense-preserving harmonic mapping f in the unit disk is defined by

$$\|S_f\| = \sup_{z\in\mathbb{D}} |S_f(z)|\cdot(1-|z|^2)^2\,.$$

It is easy to check (using the chain rule again and the Schwarz-Pick lemma) that $\|S_{f\circ\sigma}\| = \|S_f\|$ for any automorphism of the unit disk σ. For further properties of S_f and the motivation for this definition, see [5].

1.2. **An affine and linear invariant family.** In [10, 11] Pommerenke studied the so-called *linear invariant families*; that is, families of locally univalent holomorphic functions φ in the unit disk normalized by the conditions $\varphi(0) = 1 - \varphi'(0) = 0$ and which are closed under the transformation

$$\Phi_\zeta(z) = \frac{\varphi\left(\dfrac{\zeta+z}{1+\overline{\zeta}z}\right) - \varphi(\zeta)}{(1-|\zeta|^2)\varphi'(\zeta)}\,, \quad \zeta\in\mathbb{D}.$$

Let $\mathcal{F}$ be a family of sense-preserving harmonic mappings $f = h + \overline{g}$ in $\mathbb{D}$, normalized with $h(0) = g(0) = 0$ and $h'(0) = 1$. The family is said to be *affine and linear invariant* if it closed under the two operations of *Koebe transform* and *affine change*:

$$K_\zeta(f)(z) = \frac{f\left(\dfrac{z+\zeta}{1+\overline{\zeta}z}\right) - f(\zeta)}{(1-|\zeta|^2)h'(\zeta)}\,, \quad |\zeta|<1\,,$$

and

$$A_\varepsilon(f)(z) = \frac{f(z) - \overline{\varepsilon f(z)}}{1-\overline{\varepsilon}g'(0)}\,, \quad |\varepsilon|<1\,.$$

We refer to the reader to the paper [12] where Sheil-Small offers an in depth study of affine and linear invariant families $\mathcal{F}$ of harmonic mappings in $\mathbb{D}$.

Using that the Schwarzian derivative for harmonic mappings satisfies the chain rule and is invariant under affine changes $af + b\overline{f}$, $|a| \neq |b|$,

it is easy to show that the family $\mathcal{F}_\lambda$ of sense-preserving harmonic mappings $f = h+\overline{g}$ in $\mathbb{D}$ with $h(0) = g(0) = 0, h'(0) = 1$, and $||S_f|| \leq \lambda$ is affine and linear invariant. We let $\mathcal{F}_\lambda^0 = \{f \in \mathcal{F}_\lambda \colon g'(0) = 0\}$. These two families are studied in [2], where it is shown in particular that $\mathcal{F}_\lambda^0$ is a compact family of harmonic mappings with respect to the topology of uniform convergence on compact subsets of $\mathbb{D}$. In that paper, the following notation was used: an analytic function in the unit disk ω with $\omega(\mathbb{D}) \subset \mathbb{D}$ is said to belong to $\mathcal{A}_\lambda^0$ (*resp.* $\mathcal{A}_\lambda$) if there exists a harmonic mapping $f = h + \overline{g} \in \mathcal{F}_\lambda^0$ (*resp.* $\mathcal{F}_\lambda$) with dilatation ω. The quantity

$$R_\lambda = \max_{\omega \in \mathcal{A}_\lambda^0} |\omega'(0)| = \sup_{\omega \in \mathcal{A}_\lambda} \|\omega^*\|\,,$$

where

$$\|\omega^*\| = \sup_{z \in \mathbb{D}} \frac{|\omega'(z)| \cdot (1 - |z|^2)}{1 - |\omega(z)|^2}\,,$$

was shown to play a distinguished role in the analysis offered in [2].

Perhaps at this point we should mention that according to the first of the properties for the Schwarzian derivative mentioned in the previous section, we have that the family $\mathcal{F}_0$ consists only of functions of the form $f = \alpha T + \beta\overline{T}$, where $|\alpha| > |\beta|$ and T is a Möbius transformation. Therefore, all dilatations in $\mathcal{A}_0$ are constant functions.

2. Main Results

The following lemma will be important for our purposes.

Lemma 1. *As before, let* $R_\lambda = \max_{\omega \in \mathcal{A}_\lambda^0} |\omega'(0)|$. *Then*

$$\lim_{\lambda \to 0^+} R_\lambda = 0\,.$$

Proof. Since $\mathcal{F}_{\lambda_1}^0 \subset \mathcal{F}_{\lambda_2}^0$ whenever $0 < l_1 < l_2$, we have $0 \leq R_{\lambda_1} \leq R_{\lambda_2}$ as well. Therefore, we conclude that there exists $\lim_{\lambda \to 0^+} R_\lambda$ and it remains to check that this limit equals 0.

Consider an arbitrary positive number λ. By the definition of R_λ and the compacity of $\mathcal{F}_\lambda^0$, we see that for each such λ there is a harmonic mapping $f_\lambda \in \mathcal{F}_\lambda^0$ with dilatation ω_λ satisfying $|\omega_\lambda'(0)| = R_\lambda$. Since for a given $\rho > 0$ the family $\{f_\lambda \colon \lambda \leq \rho\} \subset \mathcal{F}_\rho^0$ and $\mathcal{F}_\rho^0$ is compact, we see that there is a function $f_0 \in \cap_{\rho>0}\mathcal{F}_\rho^0$ with dilatation ω_0 such that $f_\lambda \to f_0$ as $l \to 0$ uniformly on compact subsets in the unit disk (hence $\omega_\lambda'(0) \to \omega_0'(0)$ as $l \to 0$ too). Obviously, $\cap_{\rho>0}\mathcal{F}_\rho^0 = \mathcal{F}_0^0$ and the dilatations of functions in $\mathcal{F}_0^0$ are constants, thus $0 = \omega_0'(0) = \lim_{\lambda \to 0} R_\lambda$. □

We now state and prove the main theorems in this paper.

THEOREM 1. *There exists $\delta_0 > 0$ such that if $\|S_f\| \leq \delta_0$, then f is univalent.*

Proof. For any real number $\lambda > 0$, we have that if $f = h + \overline{g} \in \mathcal{F}_\lambda$, the Schwarzian norm of h is bounded by [5, Thm. 6]. Hence (see [10]),

$$\sup_{z\in\mathbb{D}} \left| \frac{h''(z)}{h'(z)} \right| (1 - |z|^2) \leq K_1$$

for some constant $K_1 > 0$. Moreover, by using ω to denote the dilatation of f, we have (see, for instance, [4]) that there exists another positive constant K_2 such that

$$\sup_{z\in\mathbb{D}} \frac{|\omega''(z)|\ (1 - |z|^2)^2}{1 - |\omega(z)|^2} \leq K_2 \|\omega^*\| \leq K_2 R_\lambda \,.$$

Hence, using (2) and the triangle inequality, we see that for any such function $f = h + \overline{g}$,

$$\|S(h)\| \leq \lambda + K_1 R_\lambda + K_2 R_\lambda + \frac{3}{2} R_\lambda^2 \,. \tag{3}$$

Now, using Lemma 1 and the fact that R_λ increases with λ, we have that there exists a unique solution δ_0, say, of the equation

$$\lambda + K_1 R_\lambda + K_2 R_\lambda + \frac{3}{2} R_\lambda^2 = 2 \,.$$

This implies by (3) that if $\lambda \leq \delta_0$, then $\|S(h)\| \leq 2$. In other words, the analytic part h of any function $f = h + \overline{g} \in \mathcal{F}_{\delta_0}$ is univalent by the classical Nehari criterion of univalence.

To prove that not only h but the function $f = h + \overline{g}$ itself is univalent whenever $f \in \mathcal{F}_{\delta_0}$, we proceed as follows. By the affine invariance property of F_{δ_0}, we see that for any $a \in \mathbb{D}$ the function $f_a = f + \overline{af}$ belongs to F_{δ_0} as well. It is easy to check that if $f_a = h_a + \overline{g_a}$, then $h_a = h + \overline{a}g$. Thus, the functions $h + \overline{a}g$ are also univalent for all $|a| < 1$. Since f is sense-preserving, an application of Hurwitz theorem gives that $h + \overline{a}g$ is indeed univalent for all $|a| \leq 1$. A direct application of [5, Prop. 2.1] shows that the function $f = h + \overline{g}$ is univalent, as was to be shown. □

We would like to point out that by finding an upper bound for the quantity R_λ in terms of λ, one could give an estimate of the value δ_0 in the previous theorem. Unfortunately, so far we are not able to obtain such upper bound.

The next result is related to a criterion for quasiconformal extension of harmonic mappings in terms of their Schwarzian norm.

THEOREM 2. *Let f be a sense-preserving harmonic mapping in the unit disk with $\|S_f\| \leq \delta_0 t$ for some $t < 1$, where δ_0 is as in Theorem 1. Assume, in addition, that the dilatation ω_f of f satisfies*

$$\|\omega_f\|_\infty = \sup_{z\in\mathbb{D}} |\omega_f(z)| < 1\,.$$

Then f can be extended to a quasiconformal map in $\widehat{\mathbb{C}}$.

Before proving this second theorem, we would like to stress that the hypotheses $\|\omega\|_\infty < 1$ cannot be removed as the following example shows.

EXAMPLE. *Consider the sense-preserving harmonic mapping $f = z+\overline{g}$, where g' equals the lens-map ℓ_α, $0 < \alpha \leq 1$, defined by*

$$\ell_\alpha(z) = \frac{\ell(z)^\alpha - 1}{\ell(z)^\alpha + 1}\,, \quad z \in \mathbb{D}\,,$$

with $\ell(z) = (1+z)/(1-z)$. Note that ℓ_1 equals the identity in the unit disk and that $\|\ell_\alpha\|_\infty = 1$ for all $0 < \alpha \leq 1$. In [7], it is explicitly checked that $\|\ell^*_\alpha\| = \alpha$.

Bearing in mind (2), that the dilatation of f is ℓ_α, and that

$$\sup_{z\in\mathbb{D}} \frac{|\ell''_\alpha(z)|\,(1-|z|^2)^2}{1-|\ell_\alpha(z)|^2} \leq K_2\|\ell^*_\alpha\| = K_2\alpha$$

for some absolute constant K_2, we have

$$\begin{aligned}
\|S_f\| &= \sup_{z\in\mathbb{D}} \left| \frac{\overline{\ell_\alpha(z)}\,\ell''_\alpha(z)}{1-|\ell_\alpha(z)|^2} + \frac{3}{2}\left(\frac{\overline{\ell_\alpha(z)}\,\ell'_\alpha(z)}{1-|\ell_\alpha(z)|^2}\right)^2 \right| (1-|z|^2)^2 \\
&\leq \sup_{z\in\mathbb{D}} \frac{|\ell''_\alpha(z)|\,(1-|z|^2)^2}{1-|\ell_\alpha(z)|^2} + \frac{3}{2}\sup_{z\in\mathbb{D}} \left| \frac{\ell'_\alpha(z)\,(1-|z|^2)}{1-|\ell_\alpha(z)|^2} \right|^2 \\
&\leq K_2\alpha + \frac{3}{2}\alpha^2\,.
\end{aligned}$$

Therefore, by choosing any α small enough, we obtain $\|S_f\| \leq \delta_0 t$ for any given $0 < t < 1$. On the other hand, the function f is not quasiconformal since its (sencond complex) dilatation coincides with ℓ_α and $\|\ell_\alpha\|_\infty = 1$.

We now prove Theorem 2.

Proof. Since we are assuming that $\|S_f\| \leq \delta_0 t$ for some $t < 1$, we have that $f \in \mathcal{F}_{\delta_0 t}$. By arguing as in the proof of the previous theorem and

using again that if $\lambda_1 \leq \lambda_2$ then $R_{\lambda_1} \leq R_{\lambda_2}$, we get

$$\begin{aligned} \|S(h)\| &\leq \delta_0 t + K_1 R_{\delta_0 t} + K_2 R_{\delta_0 t} + \frac{3}{2} R_{\delta_0 t}^2 \\ &\leq \delta_0 t + K_1 R_{\delta_0} + K_2 R_{\delta_0} + \frac{3}{2} R_{\delta_0}^2 \\ &< \delta_0 + K_1 R_{\delta_0} + K_2 R_{\delta_0} + \frac{3}{2} R_{\delta_0}^2 = 2\,, \end{aligned}$$

so that $\|S(h)\| \leq 2s$ for some $s < 1$. This shows (by the Ahlfors-Weill theorem) that the analytic part h of every function $f = h + \overline{g}$ in the family $\mathcal{F}_{\delta_0 t}$ can be extended to a K_s-quasiconformal function in $\widehat{\mathbb{C}}$, where $K_s = (1+s)/(1-s)$. Using again that the family $\mathcal{F}_{\delta_0 t}$ is invariant under affine transformations, we get that not only h but $h + ag$ (where $a \in \overline{\mathbb{D}}$) has a K_s-quasiconformal extension to $\widehat{\mathbb{C}}$. By arguing as in the proof of [6, Thm. 2], we conclude that f itself has a K-quasiconformal extension for an appropriate value of K. □

References

[1] L. V. Ahlfors and G. Weill, A uniqueness theorem for Beltrami equations, *Proc. Amer. Math. Soc.* **13** (1962), 975–978.

[2] M. Chuaqui, R. Hernández, and M. J. Martín, Affine and linear invariant families of harmonic mappings, arXiv:1405.5106 [math.CV].

[3] P. L. Duren, *Harmonic Mappings in the Plane*, Cambridge University Press, Cambridge, 2004.

[4] P. Ghatage and D. Zheng, Hyperbolic derivatives and generalized Schwarz-Pick estimates, *Proc. Amer. Math. Soc.* **132** (2004), 3309–3318.

[5] R. Hernández and M. J. Martín, Pre-Schwarzian and Schwarzian derivatives of harmonic mappings, *J. Geom. Anal.* DOI 10.1007/s12220-013-9413-x. Published electronically on April 13th, 2013.

[6] R. Hernández and M. J. Martín, Quasi-conformal extersions of harmonic mappings in the plane, *Ann. Acad. Sci. Fenn. Ser. A. I Math.* **38** (2013), 617–630.

[7] T. Hosokawa and S. Ohno, Topological structures of the set of composition operators on the Bloch space, *J. Math. Anal. Appl.* **314** (2006), 736–748.

[8] H. Lewy, On the non-vanishing of the Jacobian in certain one-to-one mappings, *Bull. Amer. Math. Soc.* **42** (1936), 689–692.

[9] Z. Nehari, The Schwarzian derivative and schlicht functions, *Bull. Amer. Math. Soc.* **55** (1949), 545–551.

[10] Ch. Pommerenke, Linear-invariante Familien analytischer Funktionen I, *Math. Ann.* **155** (1964), 108–154.

[11] Ch. Pommerenke, Linear-invariante Familien analytischer Funktionen II, *Math. Ann.* **156** (1964), 226–262.

[12] T. Sheil-Small, Constants for planar harmonic mappings, *J. London Math. Soc.* **42** (1990), 237–248.

Facultad de Ingeniería y Ciencias, Universidad Adolfo Ibáñez, Av. Padre Hurtado 750, Viña del Mar, Chile.
E-mail address: rodrigo.hernandez@uai.cl

Department of Physics and Mathematics, University of Eastern Finland, P.O. Box 111, FI-80101 Joensuu, Finland.
E-mail address: maria.martin@uef.fi